

The influence of an intramolecular hydrogen bond on the 1,3-N,S-coordination of crown ether-containing N-phosphorylthiourea with Ni II

Safin D., Babashkina M., Klein A., Sokolov F., Baranov S., Pape T., Hahn F., Krivolapov D.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Reaction of the lithium salt of N-phosphorylated thiourea [4'-benzo-15-crown-5]NHC(S)NHP(O)(OiPr)₂ (HLI) with NiII leads to the chelate complex [NiLI 2]. The metal center is found in a square-planar N₂S₂ environment formed by the CS sulfur atoms and the P-N nitrogen atoms of two deprotonated LI ligands. Reaction of [NiLI 2] and its non-crown ether-containing analog Ni[PhNHC(S)NP(O)(OiPr)₂]₂ ([NiLII 2]) with 2,2'-bipyridine (bipy) and 1,10-phenanthroline (phen) leads to the [Ni(bipy)LI,II 2] and [Ni(phen)LI,II 2] heteroligand complexes. The coordination mode is preserved for the phosphorus-containing ligands. The extraction properties of the crown ether-containing compounds towards alkali metal picrates were investigated. The molecular structures of HLI, [NiLI 2] and [Ni(bipy)LII 2] were elucidated by X-ray diffraction. © 2009 The Royal Society of Chemistry and the Centre National de la Recherche Scientifique.

<http://dx.doi.org/10.1039/b9nj00252a>
